arXiv:1212.6495v1 [math.RA] 28 Dec 2012

ON THE COSET STRUCTURE OF DISTRIBUTIVE SKEW LATTICES

JOÃO PITA COSTA[*]

Inštitut Jozef Štefan,
Jamova Cesta 39, 1000 Ljubljana, Slovenia
joao.pitacosta@ijs.si

January 1, 2013

Abstract

In the latest developments in the theory of skew lattices, the study of distributivity has been one of the main topics. The largest classes of examples of such algebras are distributive. Unlike what happens in lattices, the properties of cancellation and distributivity are independent for skew lattices. In this paper we will discuss several aspects of distributivity in the absence of commutativity, review the recent results by Kinyon and Leech on these matters and have an insight on the coset structure of those algebras that satisfy this property. We will also discuss combinatorial implications of these results.

1 Introduction

The study of skew lattices provides two perspectives that complement each other: one perspective considers skew lattices as noncommutative lattices and the other perspective sees these algebras as double bands. Due to this, many motivations are found in the intersection between lattice theory and semigroup theory. A natural partial order can be introduced maintaining several of the classical results known to lattice theory. Green's relations and other semigroup theoretical notions show an important role in the study of skew lattices.

In the recent developments in the theory of skew lattices, the study of distributivity has been one of the main topics [16]. Unlike what happens in lattices, the properties of cancellation and distributivity are independent for skew lattices. In the past two years a

---

[*] The author thanks the support of Fundação para a Ciência e Tecnologia, with the reference SFRH/BD/36694/2007, that permitted this research work during 2008-2012.

group of mathematicians have worked on this topic (namely K. Cvetko-Vah, J. Leech, M. Spinks and M. Kinyon). A skew lattice is *linearly distributive* if all its sub skew chains are distributive. Linear distributivity has been a fundamental tool in the recent study of distributive skew lattices in [10] being a necessary but not sufficient condition to distributivity. Both linear distributivity and quasi-distributivity are necessary but not sufficient conditions for distributivity. Symmetry seems to be the context where these properties come together. A nine-element example of a linearly distributive, quasi-distributive skew lattice that fails to be distributive (and also symmetric) is to appear in [10]. Though, in the presence of symmetry distributive skew lattices are those that are both quasi-distributive and linearly distributive.

The coset structure of a skew lattice permits us to derive certain coset laws characterizing the varieties of cancellative, strongly symmetric, symmetric, and categorical skew lattices. Moreover, the coset laws for distributive skew lattices are a recent achievement due to the research in [10]. From the stated coset laws we are able to derive some interesting combinatorial properties. This study started in [15] to be continued in [6], [2] and in [5]. It was our aim to complete the characterization of the subvarieties of skew lattices mentioned, having a clear overview of this study within the adopted approach.

2 Preliminaires

A skew lattice is a set S with binary operations $\wedge$ and $\vee$ that are both idempotent and associative, satisfying the absorption laws $x \wedge (x \vee y) = x = (y \vee x) \wedge x$ and their duals. A *band* is a semigroup of idempotents. A *semilattice* is a commutative band. When $\mathbf{S}$ is a commutative semigroup, the set $E(S)$ of all idempotents in $\mathbf{S}$ is a semilattice under the semigroup multiplication. When $\mathbf{S}$ is not commutative, $E(S)$ needs not be closed under multiplication [8] . Recall that a band is regular if it satisfies the identity $xyxzx = xyzx$. Skew lattices can be seen as double regular bands by considering that the band reducts $(S, \wedge)$ and $(S, \vee)$ are regular. Green's relations are five equivalence relations, introduced in [7] , characterizing the elements of a semigroup in terms of the principal ideals they generate. Due to the absorption dualities, the Green's relations in the context of skew lattices are defined in [12] by $\mathcal{R} = \mathcal{R}_\wedge = \mathcal{L}_\vee$, $\mathcal{L} = \mathcal{L}_\wedge = \mathcal{R}_\vee$ and $\mathcal{D} = \mathcal{D}_\wedge = \mathcal{D}_\vee$. Right-handed skew lattices are the skew lattices for which $\mathcal{R} = \mathcal{D}$ while left-handed skew lattices are determined by $\mathcal{L} = \mathcal{D}$ [16] . Influenced by the natural quasiorders defined in bands [8] , we define in a skew lattice $\mathbf{S}$ the following distinct concepts of order: the *natural partial order* defined by $x \geq y$ if $x \wedge y = y = y \wedge x$ or, dually, $x \vee y = x = y \vee x$; the *natural preorder* defined by $x \succeq y$ if $y \wedge x \wedge y = y$ or, dually, $x \vee y \vee x = x$. Observe that $x \,\mathcal{D}\, y$ iff $x \succeq y$ and $y \succeq x$. Usually $\mathcal{D}$ is referred in the available literature as the *natural equivalence.* The fact that $\mathcal{D}$ can be expressed by the natural preorder $\preceq$ allows us to draw diagrams that are capable of representing skew lattices as the one in Figure 1. An *admissible Hasse diagram* of (a subset of) a skew lattice is a Hasse diagram for the natural partial order (usually represented by full edges) together with an indication of all $\mathcal{D}$-congruent elements (usually represented by dashed edges).



$\wedge$	0	a	b	c	1
0	0	0	0	0	0
a	0	a	b	0	a
b	0	a	b	0	b
c	0	0	0	c	c
1	0	a	b	c	1

$\vee$	0	a	b	c	1
0	0	a	b	c	1
a	a	a	a	1	1
b	b	b	b	1	1
c	c	1	1	c	1
1	1	1	1	1	1

Figure 1: *The Caley tables and admissible Hasse diagram of the righ-handed skew lattice* $\mathbf{NC}_5^R$.

A skew lattice $\mathbf{S}$ is rectangular if $x \wedge y = y \vee x$ holds. Rectangular skew lattices may then be viewed as the conjugation of its pair of rectangular band reducts $(S, \wedge)$ and $(S, \vee)$ whose binary operations dualize each other. Moreover, whenever $\mathbf{S}$ is a skew lattice $\mathcal{D}$ is a congruence, $\mathbf{S}/\mathcal{D}$ is the maximal lattice image of $\mathbf{S}$ and all congruence classes of $\mathcal{D}$ are maximal rectangular skew lattices in $\mathbf{S}$ Recall that a *chain* (or *totally ordered set*) is a set where each two elements are (order) related, and an *antichain* is a set where no two elements are (order) related. We call $\mathbf{S}$ a *skew chain* whenever $S/\mathcal{D}$ is a chain. All $\mathcal{D}$-classes are antichains. A *primitive* skew lattice is a skew lattice composed by two comparable $\mathcal{D}$-classes, and a *diamond* (or *skew diamond*) is a skew lattice composed by two incomparable $\mathcal{D}$-classes, A and B, a join class $J = A \vee B$ and a meet class $M = A \wedge B$. In particular, we express the primitive skew lattice as $\{\, A > B \,\}$ and the skew diamond $\mathbf{X}$ as $\{\, J > A, B > M \,\}$. Note that if A and B are non-comparable $\mathcal{D}$-classes in a skew lattice $\mathbf{S}$, with $A \wedge B = M$ and $A \vee B = J$, then the $\mathcal{D}$-relation on the skew diamond $\mathbf{X} = \{\, J > A, B > M \,\}$ is the restriction of the $\mathcal{D}$-relation on $\mathbf{S}$, and the $\mathcal{D}$-classes in the sub skew lattice $\mathbf{X}$ are exactly A, B, M and J. In this case $\{\, J > A \,\}$, $\{\, J > B \,\}$, $\{\, A > M \,\}$ and $\{\, B > M \,\}$ are primitive skew lattices.

Proposition 1. *[12] Let A and B be comparable $\mathcal{D}$-classes in a skew lattice $\mathbf{S}$ such that $A \geq B$. Then, for each $a \in A$, there exists $b \in B$ such that $a \geq b$, and dually, for each $b \in B$, there exists $a \in A$ such that $a \geq b$.*

Proposition 2. *[12] Let* $\{\, J > A, B > M \,\}$ *be a skew diamond. Then, for every* $a \in A$ *there exists* $b \in B$ *such that* $a \vee b = b \vee a$ *in* J *and* $a \wedge b = b \wedge a$ *in* M*. Moreover,*

$$J = \{a \vee b \,|\, a \in A, b \in B \text{ and } a \vee b = b \vee a\} \text{ and } M = \{\, a \wedge b \mid a \in A, b \in B \text{ and } a \wedge b = b \wedge a \,\}.$$

Consider a skew lattice $\mathbf{S}$ consisting of exactly two $\mathcal{D}$-classes $A > B$. Given $b \in B$, the subset $A \wedge b \wedge A = \{a \wedge b \wedge a \,|\, a \in A\}$ of B is said to be a *coset* of A in B (or an A-*coset in* B). Similarly, a coset of B in A (or a B-coset in A) is any subset $B \vee a \vee B = \{b \vee a \vee b \,|\, b \in B\}$ of A, for a fixed $a \in A$. On the other hand, given $a \in A$, the *image set* of a in B is the set

$$a \wedge B \wedge a = \{\, a \wedge b \wedge a \mid b \in B \,\} = \{\, b \in B \mid b < a \,\}.$$

Dually, given $b \in B$ the set $b \vee A \vee b = \{\, a \in A : b < a \,\}$ is the image set of b in A (cf. [15], [17], [20] and [19]).

Theorem 3. *[15] Let* $\mathbf{S}$ *be a skew lattice with comparable* $\mathcal{D}$*-classes* $A > B$*. Then,* B *is partitioned by the cosets of* A *in* B *and the image set of any element* $a \in A$ *in* B *is a transversal of the cosets of* A *in* B*; dual remarks hold for any* $b \in B$ *and the cosets of* B *in* A *that determine a partition of* A*. Moreover, any coset* $B \vee a \vee B$ *of* B *in* A *is isomorphic to any coset* $A \wedge b \wedge A$ *of* A *in* B *under a natural bijection* φ *defined implicitly for any* $a \in A$ *and* $b \in B$ *by:* $x \in B \vee a \vee B$ *corresponds to* $y \in A \wedge b \wedge A$ *if and only if* $x \geq y$*. Furthermore, the operations* $\wedge$ *and* $\vee$ *on* $A \cup B$ *are determined jointly by the coset bijections and the rectangular structure of each* $\mathcal{D}$*-class.*

In fact, all cosets and all image sets are rectangular sub skew lattices while all coset bijections are isomorphisms between cosets (cf. [20]). Moreover, coset equality is given by the following result:

Proposition 4. *[5] Let* $\mathbf{S}$ *be a skew lattice with comparable* $\mathcal{D}$*-classes* $A > B$ *and let* $y, y' \in B$*. The following are equivalent:*

(i) $A \wedge y \wedge A = A \wedge y' \wedge A$*;*

(ii) for all $x \in A$*,* $x \wedge y \wedge x = x \wedge y' \wedge x$*;*

(iii) there exists $x \in A$ *such that* $x \wedge y \wedge x = x \wedge y' \wedge x$*.*

Dual results hold, having a similar statement.

A skew lattice is said to be *symmetric* if it is biconditionally commutative, that is if it satisfies $x \wedge y = y \wedge x$ iff $x \vee y = y \vee x$. All skew lattices for which $S/\mathcal{D}$ is a chain, are symmetric (cf. [14]). A skew lattice $\mathbf{S}$ is said to be *normal* if it satisfies $x \wedge y \wedge z \wedge w = x \wedge z \wedge y \wedge w$ and, dually, if it satisfies $x \vee y \vee z \vee w = x \vee z \vee y \vee w$ is named *conormal.* Skew lattices satisfying $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ and $(x \vee y) \wedge z = (x \wedge z) \vee (y \wedge z)$ are called $\wedge$-*distributive.* Finally, a skew lattice $\mathbf{S}$ is called *cancellative* if for all $x, y, z \in S$,

$z \vee x = z \vee y$ and $z \wedge x = z \wedge z$ imply $x = y$, and $x \vee z = y \vee z$ and $x \wedge z = y \wedge z$ imply $x = y$. In general, a *skew Boolean algebra* is an algebra $\mathbf{S} = (S; \vee, \wedge, \backslash, 0)$ such that $(S; \vee, \wedge, 0)$ is a $\wedge$- distributive, normal and symmetric skew lattice with a constant 0, named zero, satisfying $x \wedge 0 = 0 = 0 \wedge x$, and a binary operation $\backslash$ on $\mathbf{S}$, named relative complement, satisfying $(x \wedge y \wedge x) \vee (x \backslash y) = x$ and $(x \wedge y \wedge x) \wedge (x \backslash y) = 0$. Skew Boolean algebras form a variety of distributive, normal and symmetric skew lattices (cf. [13]).

Proposition 5. *[15] Let* $\mathbf{S}$ *be a skew lattice.* $\mathbf{S}$ *is symmetric if and only if given any skew diamond* $\{ J > A, B > M \}$ *in* $\mathbf{S}$ *and any* $m, m' \in M$, $j, j' \in J$ *the following equivalences hold:*

(a) $J \wedge m \wedge J = J \wedge m' \wedge J$ *if and only if* $A \wedge m \wedge A = A \wedge m' \wedge A$ *and* $B \wedge m \wedge B = B \wedge m' \wedge B$*;*

(b) $M \vee j \vee M = M \vee j' \vee M$ *if and only if* $A \vee j \vee A = A \vee j' \vee A$ *and* $B \vee j \vee B = B \vee j' \vee B$*.*

Proposition 6. *[18] Let* $\mathbf{S}$ *be a skew lattice. Then,* $\mathbf{S}$ *is normal iff for each comparable pair of* $\mathcal{D}$*-classes* $A > B$ *in* $\mathbf{S}$ *and all* $x, x' \in B$, $A \wedge x \wedge A = A \wedge x' \wedge A$*. Dually,* $\mathbf{S}$ *is conormal iff for all comparable pairs of* $\mathcal{D}$*-classes* $A > B$ *in* $\mathbf{S}$ *and all* $x, x' \in A$, $B \vee x \vee B = B \vee x' \vee B$*.*

Corollary 7. *Let* $\mathbf{S}$ *be a skew lattice. Then, for each comparable pair of* $\mathcal{D}$*-classes* $A > B$ *in* $\mathbf{S}$ *and for every* $x \in A$ *there exists a unique* $y \in B$ *such that* $y \leq x$*.*

The *variety of distributive skew lattices* was first introduced in [12] by the following defining identities (equivalent in the presence of symmetry):

$(d1)$ $x \wedge (y \vee z) \wedge x = (x \wedge y \wedge x) \vee (x \wedge z \wedge x)$.

$(d2)$ $x \vee (y \wedge z) \vee x = (x \vee y \vee x) \wedge (x \vee z \vee x)$.

Clearly, $\mathbf{S}/\mathcal{D}$ is a distributive lattice. To skew lattices as such we call *quasi-distributive.* Moreover, in the presence of normality, distributivity is equivalent to quasi-distributivity (cf. [16]). In order to explore a bit more the different concepts of distributivity available in the literature, consider the following axioms:

$(d3)$ $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$.

$(d4)$ $(x \vee y) \wedge z = (x \wedge z) \vee (y \wedge z)$.

$(d5)$ $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$.

$(d6)$ $(x \wedge y) \vee z = (x \vee z) \wedge (y \vee z)$.

Skew lattices satisfying $(d3)$ and $(d4)$ are called $\wedge$*-distributive* while skew lattices satisfying $(d5)$ and $(d6)$ are called $\vee$*-distributive.* Examples exist of skew lattices satisfying any combination of these four distributive identities (cf. [11]).

Proposition 8. *Let* $\mathbf{S}$ *be a skew lattice. Then* $\mathbf{S}$ *is distributive if and only if for all* $a, b, c \in S$*,*

(i) $a \wedge (c \vee b \vee c) \wedge a = (a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a)$ *or, dually,*

(ii) $a \vee (c \wedge b \wedge c) \vee a = (a \vee c \vee a) \wedge (a \vee b \vee a) \wedge (a \vee c \vee a)$*.*

Proof. While the direct implication is clear, the converse follows from the fact that, $(a \wedge b \wedge a) \vee (a \wedge c \wedge a) \vee (a \wedge b \wedge a)$ and $(a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a)$ are $\mathcal{D}$-related. For the details observe that, for all $x, y \in S$, $x \vee y = (y \vee x \vee y) \wedge (x \vee y \vee x)$ due to the rectangularity of $\mathcal{D}_{x \vee y \vee x}$ where both $x \vee y \vee x$ and $y \vee x \vee y$ belong. Analogously, $x \wedge y = (y \wedge x \wedge y) \vee (x \wedge y \wedge x)$, for all $x, y \in S$. Hence,

$$\begin{aligned}
&a \wedge (b \vee c) \wedge a = a \wedge (c \vee b \vee c) \wedge (b \vee c \vee b) \wedge a \\
&= a \wedge (c \vee b \vee c) \wedge a \wedge a \wedge (b \vee c \vee b) \wedge a \\
&= [(a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a)] \wedge [(a \wedge b \wedge a) \vee (a \wedge c \wedge a) \vee (a \wedge b \wedge a)] \\
&= [(a \wedge b \wedge a) \vee (a \wedge c \wedge a) \vee (a \wedge b \wedge a)] \vee [(a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a)] \\
&= (a \wedge b \wedge a) \vee (a \wedge c \wedge a)
\end{aligned}$$

as required. □

A skew lattice satisfies all four distributive laws $(d3)$ to $(d6)$ if and only if it is the direct product of a rectangular skew lattice with a distributive lattice (cf. [11]). In [14] distributive skew lattices are generalized to *bidistributive* skew lattices, a variety defined by the following identities:

$(d7)$ $x \wedge (y \vee z) \wedge w = (x \wedge y \wedge w) \vee (x \wedge z \wedge w)$.

$(d8)$ $x \vee (y \wedge z) \vee w = (x \vee y \vee w) \wedge (x \vee z \vee w)$.

Skew lattices satisfying $(d7)$ are named $\wedge$-bidistributive skew lattices while skew lattices satisfying $(d8)$ are named $\vee$-bidistributive skew lattices. Both of these classes of skew lattices constitute varieties. Moreover, $(d3)$ and $(d4)$ hold in $\mathbf{S}$ iff $\mathbf{S}$ is $\wedge$-bidistributive and symmetric. Moreover, $\mathbf{S}$ is $\wedge$-bidistributive iff it is normal and quasi-distributive, in which case $\mathbf{S}$ is also distributive (cf. [14]).

3 Orthogonality in the skew chain

A skew lattice is *categorical* if nonempty composites of coset bijections are coset bijections. Rectangular and primitive skew lattices, skew lattices in rings and skew Boolean algebras are examples of categorical skew lattices (cf. [15] and [1]).

Example. *A minimal example of a non categorical skew lattice is given by the 8-element left-handed skew chain given in Figure 2. In this example, considering the skew chain* $\{0, 4\} > \{3, 6, 1, 7\} > \{2, 5\}$*, the coset bijections are the following:*



$\wedge$	0	1	2	3	4	5	6	7
0	0	3	2	3	0	2	6	6
1	1	1	5	1	1	5	1	1
2	2	2	2	2	2	2	2	2
3	3	3	2	3	3	2	3	3
4	4	1	5	1	4	5	7	7
5	5	5	5	5	5	5	5	5
6	6	6	2	6	6	2	6	6
7	7	7	5	7	7	5	7	7

$\vee$	0	1	2	3	4	5	6	7
0	0	4	0	0	4	4	0	4
1	0	1	6	3	4	1	6	7
2	0	1	2	3	4	5	6	7
3	0	1	3	3	4	7	6	7
4	0	4	0	0	4	4	0	4
5	0	1	2	3	4	5	6	7
6	0	1	6	3	4	1	6	7
7	0	1	3	3	4	7	6	7

Figure 2: *The admissible Hasse diagram of a left-handed non categorical skew lattice.*

$$\varphi_1 : \{\,0,4\,\} \to \{\,3,1\,\},\ \varphi_2 : \{\,0,4\,\} \to \{\,6,7\,\}$$
$$\psi_1 : \{\,3,7\,\} \to \{\,2,5\,\},\ \psi_2 : \{\,6,1\,\} \to \{\,2,5\,\}$$
$$\chi : \{\,0,4\,\} \to \{\,2,5\,\}$$

Observe that 0 *has no image by* $\psi_2\varphi_1$ *and that* $\chi(0) \in \{\,2,5\,\}$ *so that* $\psi_2\varphi_1 \neq \chi$. *The reader can find a detailed study of such examples in [3] and [9] where this skew lattice is named* X_2 *and its right-handed version is named* Y_2. *A family of non categorical examples where* X_2 *and* Y_2 *belong was further studied in [9].*

A categorical skew lattice is *strictly categorical* if the compositions of coset bijections between comparable $\mathcal{D}$-classes $A > B > C$ are never empty. Rectangular and primitive skew lattices, as well as skew Boolean algebras are strictly categorical skew lattices (cf. [1]). Normal and conormal skew lattices are strictly categorical (cf. [9]). In particular, sub skew lattices of strictly categorical skew lattices are also strictly categorical.

Theorem 9. *[6] A skew chain* $\mathbf{S}$ *consisting of* $\mathcal{D}$*-classes* $A > B > C$ *is categorical iff for all elements* $a \in A$, $b \in B$ *and* $c \in C$ *satisfying* $a > b > c$, *one (and hence both) of the following equivalent statements holds:*

(i) $(A \wedge b \wedge A) \cap (C \vee b \vee C) = (C \vee a \vee C) \wedge b \wedge (C \vee a \vee C)$;

(ii) $(A \wedge b \wedge A) \cap (C \vee b \vee C) = (A \wedge c \wedge A) \vee b \vee (A \wedge c \wedge A)$.

Moreover, $\mathbf{S}$ *is strictly categorical iff in addition to* (i)–(ii), *for all* $b, b' \in B$,

$$(A \wedge b \wedge A) \cap (C \vee b' \vee C) \neq \emptyset.$$



$\wedge$	0	a	b	1
0	0	0	0	0
a	0	a	b	a
b	0	a	b	b
1	0	a	b	1

$\vee$	0	a	b	1
0	0	a	b	1
a	a	a	a	1
b	b	b	b	1
1	1	1	1	1

Figure 3: The Caley tables and the admissible Hasse diagram of a non strictly categorical right-handed skew lattice.

Example. *A minimal example of a categorical skew lattice that is not strictly categorical is given by the right-handed manifestation of the skew chain with three $\mathcal{D}$-classes in Figure 3. In fact, the composition of the coset bijections $\psi : \{1\} \to \{a\}$ and $\varphi' : \{b\} \to \{0\}$ is empty. Observe that $\chi : \{1\} \to \{0\}$ can be decomposed either by the composition of ψ and $\varphi : \{a\} \to \{0\}$, or by the composition of $\psi' : \{1\} \to \{b\}$ and φ' (cf. [9]).*

The following is a practical criteria to identify strictly categorical skew lattices.

Proposition 10. *[9] A skew chain $A > B > C$ is strictly categorical if and only if given $a \in A$, $b, b' \in B$ and $c \in C$ such that $a > b > c$ and $a > b' > c$, then $b = b'$ must follow.*

Remark 11. *Let $A \geq B$ be comparable $\mathcal{D}$ classes in a normal skew lattice $\mathbf{S}$ and $a \in A$. Due to the normality of $\mathbf{S}$, for all $a \in A$ there exists a unique $b \in B$ such that $a \geq b$. Observe that, considering $a \wedge b \wedge a \in B$ we get $a \geq b$ and, for any other $b' \in B$ such that $a \geq b'$, we also get*

$$b = a \wedge b \wedge a = a \wedge b \wedge b' \wedge b \wedge a = a \wedge b \wedge b' \wedge b' \wedge b \wedge a = a \wedge b' \wedge b \wedge b \wedge b' \wedge a = a \wedge b' \wedge a = b'.$$

Thus, the lower $\mathcal{D}$-class in any maximal primitive sub skew lattice of a normal skew lattice has exactly one coset. Moreover, coset bijections are closed under composition with the composition of adjacent coset bijections being nonempty. Similar remarks hold regarding the coset structure of conormal skew lattices.

Example. *Strictly categorical skew lattices need not be normal: the admissible Hasse diagram in Figure 4 represents a right-handed skew chain defined by the respective Caley*



$\wedge$	a	b	c	d	1
a	a	b	c	d	a
b	a	b	c	d	b
c	c	d	c	d	c
d	c	d	c	d	d
1	a	b	c	d	1

$\vee$	a	b	c	d	1
a	a	a	a	a	1
b	b	b	b	b	1
c	a	a	c	c	1
d	b	b	d	d	1
1	1	1	1	1	1

Figure 4: The Caley tables and the admissible Hasse diagram of a non normal but strictly categorical right-handed skew lattice.

tables. It was shown to be strictly categorical but normality fails as the upper $\mathcal{D}$-class determines more then one coset in the lower $\mathcal{D}$-class: observe that, considering $A = \{1\}$, $B = \{a, b\}$ and $C = \{c, d\}$, $A > B > C$ is a strictly categorical skew chain, according to Proposition 10, but

$$A \wedge a \wedge A = \{a\} \neq \{b\} = A \wedge b \wedge A.$$

The *natural graph* of a skew lattice $\mathbf{S}$ is the undirected graph (S, E) given by the natural partial order of $\mathbf{S}$, where S is the set of vertices and $\{x, y\}$ forms an edge in E whenever $x > y$ or $y > x$. The natural graph of a skew lattice $\mathbf{S}$ is the admissible Hasse diagram of the skew lattice without the dashed edges, representing only the partial order.

A skew lattice is *connected* when its natural graph is connected, that is, when for every pair of vertices (x, y), the graph contains a path from x to y. A *component* of $\mathbf{S}$ is any connected component of its graph. The Hasse diagram of the lattice $\mathbf{S}/\mathcal{D}$ is a connected graph. Thus, each component of $\mathbf{S}$ has a nonempty intersection with each equivalence class of $\mathbf{S}$.

Theorem 12. *[15] The maximal connected sub algebras. of a skew lattice* $\mathbf{S}$ *are its components. Moreover, the partition of* $\mathbf{S}$ *into components is a congruence partition for which the induced quotient algebra is the maximal rectangular image of* $\mathbf{S}$*.*

Given $\mathcal{D}$-classes $A \geq B$, by an *order component* of A in B is meant the intersection of a component of $A \cup B$ with B. Order components were introduced in [15] as "class components". We will name them differently to distinguish them from other types of components that arise in this dissertation. Due to Theorem 12, order components form a partition of all the components of $\mathbf{S}$. Let $\mathbf{S}$ be a skew lattice with three equivalence classes A, B and C such that B is comparable to both A and C. A and C are *orthogonal* in B if each order component of A in B is contained in a (unique) coset of C in B, and likewise each order component of C in B is contained in a (unique) coset of A in B. Orthogonality was further studied in [15].

Example. *The right-handed skew lattice represented in Figure 1 is an example of skew lattice with $\mathcal{D}$-classes $\{ 0 \}$ and $\{ 1 \}$ that are orthogonal in $\{ c \}$ but not in $\{ a, b \}$. Observe that $b \wedge \{ 1 \} = \{ b \}$ and $\{ 0 \} \vee b = \{ b \}$. On the other hand, the order components of $\{ 0 \}$ in $\{ a, b \}$ are $\{ a \}$ and $\{ b \}$ coinciding with the order components of $\{ 1 \}$ in $\{ a, b \}$. The order component of both $\{ 0 \}$ and $\{ 1 \}$ in $\{ c \}$ is clearly $\{ c \}$.*

Proposition 13. *Let $\mathbf{S}$ be a primitive skew lattice with comparable $\mathcal{D}$-classes $A > B$. Then, the order component K of A in B is the union of image sets of elements of $K' \cap A$ in B, where K' is a component of $A \cup B$. Similar remarks hold for order components of B in A.*

Proof. Let $a \in A$. Observe that $(a \wedge B \wedge a) \cup \{ a \}$ is a connected sub skew lattice and, therefore, part of a unique component K of $\mathbf{S}$ due to Theorem 12. Hence, $K \cap B$ is an order component of A in B that contains $a \wedge B \wedge a$.

Conversely, let K' be a component of $A \cup B$ and consider the order component $K = K' \cap B$. Let $x \in K$. Due to Lemma 1, there exists $a \in A$ such that $a \geq x$. Thus $a \in K'$ and $x \in a \wedge B \wedge a$ by Theorem 3 (ii). $\square$

Whenever $\mathbf{S}$ is a skew lattice with three equivalence classes A, B and C such that B is comparable to both A and C, we say that $x \in A$ is *covered* by a coset of C in B if the image set of x in B is a subset of this coset of C. The dual definition is similar.

Lemma 14. *[15] Let $\mathbf{S}$ be a skew lattice with three equivalence classes A, B and C such that B is comparable to both A and C. Then, A and C are orthogonal in B iff each $x \in A$ is covered by a coset of C in B and, dually, if each $y \in C$ is covered by a coset of A in B.*

Proposition 15. *[15] Let $\mathbf{S}$ be a skew lattice with three equivalence classes A, B and C such that B is comparable to both A and C. If A and C are orthogonal in B, then each coset of A in B has nonempty intersection with each coset of C in B. Moreover, all such coset intersections have common cardinality.*

Theorem 16. *[18] Let $\mathbf{S}$ be a skew lattice. The following statements are equivalent:*

(i) $\mathbf{S}$ *is categorical;*

(ii) for all distinct $\mathcal{D}$-classes $A > B > C$ with elements $b \in B$ and $c \in C$ that satisfy $b > c$, the coset bijection $\phi_{b,c} : C \vee b \vee C \to B \wedge c \wedge B$ restricted to the corresponding AC-coset $(A \wedge b \wedge A) \cap (C \vee b \vee C)$ in B is a bijection of the later onto the A-coset $A \wedge c \wedge A$ in C;



Figure 5: The partition diagram of $\mathcal{D}$-classes A, B, C, with A and C orthogonal in B.

(iii) for all distinct $\mathcal{D}$-classes $A > B > C$ with elements $a \in A$ and $b \in B$ that satisfy $a > b$, the coset bijection $\phi_{b,a} : A \wedge b \wedge A \to B \vee a \vee B$ restricted to the corresponding AC-coset $(A \wedge b \wedge A) \cap (C \vee b \vee C)$ in B is a bijection of the latter onto the C-coset $C \vee a \vee C$ in A.

Due to Proposition 15, we can visualize any $\mathcal{D}$-class B as in the diagram of Figure 5. As $\mathcal{D}$-classes and cosets are rectangular skew lattices due to the rectangularity of cosets we will represent them by rectangles in this diagram, Here, B is a doubly partitioned rectangle arising from a horizontal partition by one class of congruent rectangles, the A-cosets, and a vertical partition by a second class of congruent rectangles, the C-cosets. To the nonempty intersections of A-cosets in B with C-cosets in B we call *AC-cosets* in B in conformity with [10]. We call a diagram representing intersection of partitions, like the one in Figure 5, a *partition diagram.*

Proposition 17. *[15] Let* $\mathbf{S}$ *be a skew lattice with two non comparable $\mathcal{D}$-classes A and B. Then, A and B are orthogonal in their meet class M; dually, they are orthogonal in their join class J. Moreover, individual joins and meets of pairs of elements from each class are determined by the orthogonality relationship.*

Given a skew diamond $\{ J > A, B > M \}$, J and M are each partitioned by their A-cosets, their B-cosets, the resulting AB-coset intersections and by the cosets of each in the other. In general, M-cosets in J [J-cosets in M] refine the partition by AB-coset intersections.

Theorem 18. *[15] A skew lattice* $\mathbf{S}$ *is symmetric if and only if for any skew diamond $\{ J > A, B > M \}$ in* $\mathbf{S}$*, the partitions that J and M induce on each other coincide with the partitions by the AB-coset intersections.*

Proposition 19. *[9] Given a strictly categorical skew chain $A > B > C$,*

(i) for any $a \in A$, there exists $b \in B$ such that $a \wedge B \wedge a \subseteq C \vee b \vee C$.

(ii) for any $c \in C$, there exists $b' \in B$ such that $c \vee B \vee c \subseteq A \wedge b' \wedge A$.

Proposition 20. *Let* **S** *be a categorical skew lattice. Then, A and C are orthogonal in B if and only if $A > B > C$ is a strictly categorical skew chain.*

Proof. Due to Proposition 15, the orthogonality of A and C in B implies that each coset of A in B has nonempty intersection with each coset of C in B. Proposition 9 then implies that the skew chain $A > B > C$ is strictly categorical.

Conversely, let $A > B > C$ be a strictly categorical skew chain and X be a order component of A in B. Then, It follows from Lemma 14 that the orthogonality of A and C in B is equivalent to the conditions (i) and (ii) of Proposition 19. Clearly, if A and C are orthogonal in B then Proposition 13 implies both (i) and (ii). Conversely, assume that (i) and (ii) hold and let K be a order component of A in B. Let $x \in A$. By the assumptions, $x \wedge B \wedge x$ lies in a unique C-coset in B, say $C \vee b \vee C$ for some $b \in B$. Let $y \in K$. Then, there exists a sequence of image sets $x_i \wedge B \wedge x_i$ with $x = x_1, x_2, \ldots, x_n$ such that $y \in x_n \wedge B \wedge x_n$, $(x_i \wedge B \wedge x_i) \cap (x_j \wedge B \wedge x_j) \neq \emptyset$ for every $1 \leq i, j \leq n$ and $K = \bigcup_{1 \leq i \leq n} (x_i \wedge B \wedge x_i)$ due to Proposition 13. Thus

$$\bigcap_{i \in I} \Big((x_i \wedge B \wedge x_i) \cap (C \vee b \vee C) \Big) \neq \emptyset$$

so that $K = \bigcup_{1 \leq i \leq n} (x_i \wedge B \wedge x_i) \subseteq (C \vee b \vee C)$, by Theorem 3(i). Hence, A and C are orthogonal in B. ☐

Corollary 21. *Let* **S** *be a skew Boolean algebra. Then, A and C are orthogonal in B, for all skew chains $A > B > C$ in* **S**.

Proof. All skew Boolean algebras are normal skew lattices by definition. Thus, they are strictly categorical skew lattices due to normality. Hence, Proposition 20 implies that in all skew chains $A > B > C$ in the skew Boolean algebra, A and C are orthogonal in B, as required. ☐

Proposition 22. *[10] Let $A > B > C$ be a strictly categorical skew chain, and $a \in A$, $c \in C$ such that $a > c$. Then, there exists a unique element $b \in B$ such that $a > b > c$. Furthermore, b lies jointly in the C-coset in B containing all images of a in B and in the A-coset in B containing all images of c in B.*

4 Coset laws for distributive skew lattices

The following results make use of identities involving cosets to characterize categorical and strictly categorical skew lattices by the description of their coset structure. These will lead us to the recent achievements by Leech and Kinyon in [10] that can be revisited in this coset structure context permitting us a similar characterization for distributive skew lattices.

Proposition 23. *[18] A skew lattice is categorical if and only if all its skew chains are categorical. Furthermore, given a skew chain* $\{A > B > C\}$ *in a skew lattice* $\mathbf{S}$*, the following statements are equivalent:*

- *(i)* $\{A > B > C\}$ *is categorical,*
- *(ii) For each* $c, c' \in C$, $A \wedge c \wedge A = A \wedge c' \wedge A$ *if and only if* $B \wedge c \wedge B = B \wedge c' \wedge B$ *and, for any* b, b' *in some* C*-coset in* B *such that* $b > c$ *and* $b' > c'$, $A \wedge b \wedge A = A \wedge b' \wedge A$ *[where* $b' = c' \vee b \vee c'$*].*
- *(iii) For each* $a, a' \in A$, $C \vee a \vee C = C \vee a' \vee C$ *if and only if* $B \vee a \vee B = B \vee a' \vee B$ *and, for any* b, b' *in some* A*-coset in* B *such that* $a > b$ *and* $a' > b'$, $C \vee b \vee C = C \vee b' \vee C$ *[where* $b' = a' \wedge b \wedge a'$*].*

Proposition 24. *[9] Given any categorical skew chain* $A > B > C$ *of* $\mathcal{D}$*-classes in* $\mathbf{S}$ *and* $a \in A$, $b \in B$ *and* $c \in C$ *such that* $a > b > c$*, the following statements are equivalent:*

- *(i)* $A > B > C$ *is strictly categorical.*
- *(ii) For each coset bijection* $\chi : C \vee a \vee C \to A \wedge c \wedge A$*, there exist unique coset bijections* $\varphi : B \vee a \vee B \to A \wedge b \wedge A$ *and* $\psi : C \vee b \vee C \to B \wedge c \wedge B$ *such that* $\chi = \psi \circ \varphi$.

Proposition 24 is a recent characterization of strictly categorical skew lattices in which Example 3 takes an important role. Moreover, a unique factorization of χ in Proposition 24 (*iii*) occurs precisely when such an algebra does not come up (cf. [10]). Furthermore, the following result extends this discussion to the particular case of strictly categorical skew lattices.

Proposition 25. *[10] Strictly categorical, quasi-distributive skew lattices are distributive, being cancellative when they are also symmetric.*

In [15], Leech gives us a good hint on what the coset law for the particular case of normal skew lattices could be. This hint is developed in the next result:

Proposition 26. *[18] Let* $\mathbf{S}$ *be a skew lattice. Then,* $\mathbf{S}$ *is normal iff for each comparable pair of* $\mathcal{D}$*-classes* $A > B$ *in* $\mathbf{S}$*,* B *is the entire coset of* A *in* B*. That is, for all* $x, x' \in B$*,*

$$A \wedge x \wedge A = A \wedge x' \wedge A.$$

Dually, $\mathbf{S}$ *is conormal iff for all comparable pairs of* $\mathcal{D}$*-classes* $A > B$ *in* $\mathbf{S}$ *and all* $x, x' \in A$*,* $B \vee x \vee B = B \vee x' \vee B$*.*

Example. *Proposition 26 implies that given any comparable* $\mathcal{D}$*-classes* $A > B$*,* B *is a single* A*-coset. All primitive skew lattices are categorical. Thus, to find an example of a categorical skew lattice that is not normal it suffices to take a primitive skew lattice* $A > B$ *such that* B *is not a single* A*-coset. For instance, take a right-handed skew lattices* $\mathbf{S}$ *on* $\mathcal{D}$*-classes* $\{ a, a' \} > \{ b, b' \}$ *such that both* b, b' *are below both* a, a'*. Then* $a \wedge b \wedge b' \wedge a = b'$ *while* $a \wedge b' \wedge b \wedge a = b$*. Hence,* $\mathbf{S}$ *is categorical but not normal.*

The following is based on the recent research by Leech and Kinyon on distributive skew lattices (cf. [9] and [10]). It unveils the relation between distributivity and categoricity, permitting us to develop the work done in Proposition 23 for categorical skew lattices and extend the study to the distributive case. Though, the coset law to be revealed is no more then the review of one of their recent results of [10].

Given a categorical skew chain $A > B > C$, two elements $b, b' \in B$ are *AC-connected* if a finite sequence $b = b_0, b_1, \ldots, b_n = b'$ exists in B such that $A \wedge b_i \wedge A = A \wedge b_{i+1} \wedge A$ or $C \vee b_i \vee C = C \vee b_{i+1} \vee C$ for all $0 \leq i \leq n-1$. If $(A \wedge b \wedge A) \cap (C \vee b' \vee C) \neq \emptyset$ for some $b, b' \in B$, then all elements in such an AC-coset are clearly AC-connected. An *AC-component* is any maximal AC-connected subset of B.

Example. *For instance, in Figure 2 the set* $\{ 3, 6, 1, 7 \}$ *is an* AC*-component of the represented skew lattice for which* $A = \{ 0, 4 \}$ *and* $C = \{ 2, 5 \}$*. On the other hand, considering the skew lattice in Figure 1 with* $A = \{ 1 \}$ *and* $C = \{ 0 \}$*, the* AC*-components in this case are* $\{ a \}$*,* $\{ b \}$ *and* $\{ c \}$*.*

Proposition 27. *Let* $A > B > C$ *be a categorical skew chain. Every* AC*-component is a union of* A*-cosets in* B *with* C*-cosets in* B*. Furthermore,* B *is a disjoint union of all its* AC*-components and each* AC*-component* B' *in* B *is a disjoint union of all the* AC*-cosets in* B'*.*

Proof. Let B' be a AC-component of B, $x \in B'$ and $y \in A \wedge x \wedge A$. Then, x and y are in the same A-coset in B and, therefore, they are AC-connected. Thus, $A \wedge x \wedge A \subseteq B'$ and, similarly, $C \vee x \vee C \subseteq B'$.

The second statement is now a direct consequence of Theorem 3 (i) due to the fact that cosets partition $\mathcal{D}$-classes and the AC-cosets are the elements of the double partition determined by the A-cosets in B together with the C-cosets in B. □

Corollary 28. *Let* $A > B > C$ *be a strictly categorical skew chain. Then,* B *has one only* AC*-component.*

Proof. This is a direct consequence of the definitions of AC-coset and AC-component, together with Proposition 27 and the fact that, in a strictly categorical skew chain, each A-coset in B intersects all C-cosets in B. $\square$

The converse of this result doesn't hold as it was shown in [10] where the authors present a categorical skew chain $A > B > C$ with one unique AC-coset in B that is not distributive (and, therefore, not strictly categorical).

Given a categorical skew chain $A > B > C$, the A-cosets in B are *amenable* to the C-cosets in B if in any AC-component of B each A-coset meets each C-coset. Amenability is of *coset nature* as it is shown in the next result.

Proposition 29. *Let $A > B > C$ be a categorical skew chain. The A-cosets in B are amenable to the C-cosets in B if and only if, in any AC-component B' of B, they satisfy the following equivalence for all $x, y \in B'$: $A \wedge x \wedge A = A \wedge u \wedge A$ and $C \vee y \vee C = C \vee u \vee C$ for some $u \in B'$ iff $A \wedge y \wedge A = A \wedge v \wedge A$ and $C \vee x \vee C = C \vee v \vee C$ for some $v \in B'$.*

Proof. Let B' be a AC component of B and $x, y \in B'$. Then, x and y are AC-connected. Assume that $A \wedge x \wedge A = A \wedge u \wedge A$ and $C \vee y \vee C = C \vee u \vee C$ for some $u \in B'$. Thus, the amenability of the A-cosets in B to the C-cosets in B implies the existence of $v \in (A \wedge y \wedge A) \cap (C \vee x \vee C)$ and, therefore, $A \wedge y \wedge A = A \wedge v \wedge A$ and $C \vee x \vee C = C \vee v \vee C$.

Conversely, let $x, y \in B$ such that x, y are AC-connected. Suppose, without loss of generality that $u \in B$ exists such that $A \wedge x \wedge A = A \wedge u \wedge A$ and $C \vee y \vee C = C \vee u \vee C$. Thus, $A \wedge y \wedge A = A \wedge v \wedge A$ and $C \vee x \vee C = C \vee v \vee C$ for some $v \in B$. As x and y are arbitrary elements in B, each A-coset meets each C-coset in the AC-component of B where x and y belong. $\square$

Proposition 30. *[10] Let $A > B > C$ be a categorical skew chain for which A-cosets and C-cosets in B are amenable. Then, in each AC-component B' of B, the following statements hold:*

- *i) for any $a \in A$, all images of a in B' lie in a unique C-coset in B';*
- *ii) for any $c \in C$, all images of c in B' lie in a unique A-coset in B'.*

Furthermore, for all $a \in A$ and $c \in C$ such that $a > c$ a unique element $b \in B'$ exists in each AC-component B' of B such that $a > b > c$. This b lies jointly in the C-coset in B' containing all images of a in B' and in the A-coset in B' containing all images of c in B'.

Proposition 30 can be proved with similar arguments to the ones used in Propositions 22 and 19. According to it, whenever A-cosets and C-cosets in B are amenable in a categorical skew chain $A > B > C$, A and C are "orthogonal" in each AC-component B' of B, in the sense of Proposition 20.

The following Lemma is not only a technical result but unveils the key for the coset approach to linearly distributive skew lattices. It's proof is a combination of Lemmas from the recent research to be published in [10]. We will revisit the proof of this Lemma and the following Theorem under the light of the coset laws for categorical skew lattices established in Proposition 23.

Lemma 31. *[10] A categorical skew chain $A > B > C$ is distributive if and only if for all $a \in A$, $b \in B$ and $c \in C$ with $a \succ b \succ c$ such that $a > c$,*

$$a \wedge (c \vee b \vee c) \wedge a = c \vee (a \wedge b \wedge a) \vee c. \tag{1}$$

Proof. Let $a \in A$, $b \in B$ and $c \in C$ with $a \succ b \succ c$ such that $a > c$. Observe that the expression $(a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a)$ reduces to $c \vee (a \wedge b \wedge a) \vee c$. Hence, due to Proposition 8, distributivity implies the identity (1).

Conversely, observe that the case $a \succ b \succ c$ is the only situation when the identity (1) is not trivial. Otherwise put, $a \wedge (c \vee b \vee c) \wedge a \neq (a \wedge b \wedge a) \vee (a \wedge c \wedge a) \vee (a \wedge b \wedge a)$ implies $a \succ b \succ c$ (cf. [10] Lemma 3.3). Let us now show that **S** is distributive, given $a \succ b \succ c$ with respective $\mathcal{D}$-classes $A > B > C$, under the assumption that the identity 1 holds whenever $a > c$. Consider $c' = a \wedge c \wedge a$. Observe that $a > c'$ and that

$$(a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a) = c' \vee (a \wedge b \wedge a) \vee c' \tag{2}$$

As $A \wedge c \wedge A = A \wedge c' \wedge A$ and **S** is categorical, Proposition 23 implies that $c \vee b \vee c$ and $c' \vee b \vee c'$ lie in the same A-coset in B so that, by Proposition 4,

$$a \wedge (c \vee b \vee c) \wedge a = a \wedge (c' \vee b \vee c') \wedge a. \tag{3}$$

Due to the assumption, $c' \vee (a \wedge b \wedge a) \vee c' = a \wedge (c' \vee b \vee c') \wedge a$. From the identities (3) and (2) then follows

$$a \wedge (c \vee b \vee c) \wedge a = (a \wedge c \wedge a) \vee (a \wedge b \wedge a) \vee (a \wedge c \wedge a)$$

and, therefore, **S** is distributive. □

Theorem 32. *[10] Given any categorical skew chain $A > B > C$ in a skew lattice **S** the following are equivalent:*

i) $A > B > C$ is distributive;

ii) For all AC components B' of B and $x, y \in B'$, $A \wedge x \wedge A = A \wedge u \wedge A$ and $C \vee y \vee C = C \vee u \vee C$ for some $u \in B$ iff $A \wedge y \wedge A = A \wedge v \wedge A$ and $C \vee x \vee C = C \vee v \vee C$ for some $v \in B$.

Proof. Assume (i) and let B' be an AC component of B, $x, y \in B'$ and $z \in (A \wedge x \wedge A) \cap (C \vee y \vee C)$ in B'. Thus, $z = a \wedge x \wedge a$ and $y = c \vee z \vee c = c \vee (a \wedge x \wedge a) \vee c$, for some $a \in A$ and $c \in C$. Due to distributivity,

$$y = (c \vee a \vee c) \wedge (c \vee x \vee c) \wedge (c \vee a \vee c).$$

Take $v = c \vee x \vee c$. Hence, $A \wedge y \wedge A = A \wedge (c \vee x \vee c) \wedge A = A \wedge v \wedge A$ and $C \vee x \vee C = C \vee (c \vee x \vee c) \vee C = C \vee v \vee C$. The proof for the converse direction of this statement is analogous.

Conversely, assume (ii) and let $a \succ b \succ c$ with corresponding $\mathcal{D}$-classes $A > B > C$ in $\mathbf{S}$ such that $a > c$. Observe that $a \wedge (c \vee b \vee c) \wedge a$ and $c \vee (a \wedge b \wedge a) \vee c$ are AC-connected by the following argument:

$$C \vee (c \vee (a \wedge b \wedge a) \vee c) \vee C = C \vee (a \wedge b \wedge a) \vee C,$$

$$A \wedge (a \wedge b \wedge a) \wedge A = A \wedge b \wedge A,$$

$$C \vee b \vee C = C \vee (c \vee b \vee c) \vee C \text{ and}$$

$$A \wedge (c \vee b \vee c) \wedge A = A \wedge (a \wedge (c \vee b \vee c) \wedge a) \wedge A.$$

Moreover, $a > a \wedge (c \vee b \vee c) \wedge a, c \vee (a \wedge b \wedge a) \vee c > c$ due to the fact that $a > c$.

Propositions 29 and 30 then imply the equality $a \wedge (c \vee b \vee c) \wedge a = c \vee (a \wedge b \wedge a) \vee c$ that, by Lemma 31, is sufficient to achieve the distributivity of $\mathbf{S}$. ☐

The above result is due to the recent research by Kinyon and Leech on distributivity for skew lattices. It shows how the property of distributivity in skew chains is of *coset nature*. The following result is a simple consequence of this, being a particular case of a more general result due to [10].

Corollary 33. *All strictly categorical skew chains are distributive.*

Proof. Given any strictly categorical skew chain $A > B > C$ in a skew lattice $\mathbf{S}$, Corollary 28 implies that B has one only AC-component, B itself. Let $x, y \in B$ and assume that $A \wedge x \wedge A = A \wedge u \wedge A$ and $C \vee y \vee C = C \vee u \vee C$ for some $u \in B$. As $A > B > C$ is strictly categorical then also $A \wedge y \wedge A$ and $C \vee x \vee C$ intersect in some $v \in B$ so that $A \wedge y \wedge A = A \wedge v \wedge A$ and $C \vee x \vee C = C \vee v \vee C$. Proposition 32 then implies the distributivity of $A > B > C$. ☐

Though, the converse statement to Corollary 33 does not hold: the skew lattice represented in Figure 3 is an example of a distributive skew lattice that is not strictly categoricalhaving two singular AC-components.

Corollary 34. *[10] Let $A > B > C$ be a skew chain. Then, $A > B > C$ is distributive if and only if $A > B' > C$ is strictly categorical, for all AC-component B' in B.*

Proof. Assume that $A > B > C$ is a distributive skew chain. Then, $A > B > C$ is categorical and so are all its sub skew lattices. Let B_0 be an AC-component of B and consider the categorical skew chain $A > B_0 > C$. Let $b, b' \in B_0$. As B_0 is an AC-component of B, b and b' are AC-connected. Hence, there exists a finite sequence $b = b_0, b_1, \ldots, b_n = b'$ in B_0 such that $A \wedge b_i \wedge A = A \wedge b_{i+1} \wedge A$ or $C \vee b_i \vee C = C \vee b_{i+1} \vee C$ for all $0 \leq i \leq n-1$. Due to Proposition 32, without loss of generality there exists $b'' \in B_0$ such that $A \wedge b \wedge A = A \wedge b'' \wedge A$ and $C \vee b'' \vee C = C \vee b' \vee C$ following that $A > B_0 > C$ is strictly categorical.

Conversely, assume that $A > B' > C$ is strictly categorical, for all AC-components B' in B. Then, each of these skew chains are distributive, due to Corollary 33. Thus, Lemma 31 and Proposition 27 ensure the distributivity of $A > B > C$. ☐

In Figure 6 is presented a diagram relating the varieties of skew lattices mentioned during this paper regarding the characterization of subvarieties of skew lattices by coset laws. These are the relationships known to date due to the woks published in [12], [13], [14], [15], [16], [2], [3], [10] and [18].

5 Combinatorial Implications

Several are the instances of combinatorial implications in the literature when reading through the study of the coset structure of a skew lattice. We will derive certain combinatorial properties that follow from the results of the coset laws. We shall do this by counting the cosets determined by a $\mathcal{D}$-class in another comparable with this. The results in this chapter were motivated by studies developed in [15] and [3] and are due to the recent research developed in [4], [5] and [18].

In this section we turn our attention to the image set of an element in a coset and its role of a transversal of cosets permitting us to count those cosets. Transversals allows us to define the *index* of B in A, first presented in [3] and denoted by $[A : B]$, as the cardinality of the image set $b \vee A \vee b$, for any $b \in B$. Dually, we define the index of A in B, denoted by $[B : A]$, as the cardinality of the image set $a \wedge B \wedge a$, for any $a \in A$. The index $[A : B]$ equals the cardinality of the set of all B-cosets in A, and $[B : A]$ equals the cardinality of the set of all A-cosets in B. As all A-cosets in B and all B-cosets in A have a common size due to coset decomposition, we name this number the *order of the A-coset in B* (or the order of the B-coset in A), denoting it by $\omega[A, B]$ or, equivalently, by $\omega[B, A]$ (denoted by $c[A, B]$ in [12]).

Example. *The situation in $A > B$ is illustrated in the following figure where the upper egggoxes represent the B-cosets in A and the lower eggboxes represent the A-cosets in B. In this case, $|A| = 18$ and $|B| = 12$ with $[A, B] = 3$, $[B, A] = 2$ and $\omega[A, B] = 6$.*

$$\begin{tikzcd}
 & & \mathbb{S}_R \arrow[d] & & \\
 & & \mathbb{S}_{MD} \arrow[dll] \arrow[dr] & & \\
\mathbb{S}_N \arrow[dd] & & & \mathbb{S}_S \cup \mathbb{S}_{C^*} \cup \mathbb{S}_{QD} \arrow[dl] \arrow[dd] \arrow[ddr] & \\
 & & \mathbb{S}_{C^*} \cup \mathbb{S}_{QD} \arrow[dll] \arrow[d] & & \\
\mathbb{S}_{C^*} \arrow[ddd] & & \mathbb{S}_D \arrow[dl] \arrow[dd] & \mathbb{S}_K \arrow[ddl] \arrow[dd] & \mathbb{S}_S \cup \mathbb{S}_C \arrow[ddl] \\
 & \mathbb{S}_{LD} \arrow[ddl] & & & \\
 & & \mathbb{S}_{QD} \arrow[dd] & \mathbb{S}_{S^*} \arrow[d] & \\
\mathbb{S}_C \arrow[drr] & & & \mathbb{S}_S \arrow[dl] & \\
 & & \mathbb{S} & &
\end{tikzcd}$$

$\mathbb{L} \equiv$ Lattices (or Commutative Skew Lattices)

$\mathbb{S} \equiv$ Skew Lattices

$\mathbb{S}_R \equiv$ Rectangular Skew Lattices

$\mathbb{S}_{MD} \equiv \mathbb{S}_S \cup \mathbb{S}_N \cup \mathbb{S}_{QD} \equiv \wedge$-distributive Skew Lattices

$\mathbb{S}_N \equiv$ Normal Skew Lattices

$\mathbb{S}_{C^*} \equiv$ Strictly Categorical Skew Lattices

$\mathbb{S}_D \equiv$ Distributive Skew Lattices

$\mathbb{S}_{LD} \equiv$ Linearly Distributive Skew Lattices

$\mathbb{S}_{QD} \equiv$ Quasi-distributive Skew Lattices

$\mathbb{S}_C \equiv$ Categorical Skew Lattices

$\mathbb{S}_K \equiv$ Cancellative Skew Lattices

$\mathbb{S}_{S^*} \equiv$ Strongly Symmetric Skew Lattices

$\mathbb{S}_S \equiv$ Symmetric Skew Lattices

Figure 6: *Diagram relating the varieties of skew lattices.*



As a direct consequence of the coset decomposition stated in Theorem 3 follows:

Lemma 35. *Given a skew lattice* $\mathbf{S}$ *with comparable* $\mathcal{D}$*-classes* $X > Y$ *consider the set of all* Y*-cosets in* X*,* $\{X_i \mid i \leq [X : Y]\}$ *and the set of* X*-cosets in* Y*,* $\{Y_j \mid j \leq [Y : X]\}$*. Then,* X *is finite if and only if* $[X : Y]$ *and* $\omega[X, Y]$ *are finite and, in that case,*

$$|X| = [X : Y].\omega[X, Y].$$

Similar remarks hold regarding the finitude of Y *and, likewise,* $|Y| = [Y : X].\omega[X, Y]$ *whenever* Y *is finite.*

Whenever $\{ J > A, B > M \}$ is a skew diamond in a skew lattice $\mathbf{S}$ and $J = \bigcup_I J_i^A = \bigcup_K J_k^B$ are the respective partitions of J into A-cosets and B-cosets, Propositions 15 and 17 imply that all possible intersections $J_i^A \cap J_k^B$ have common cardinality. In particular, they are all nonempty. Dual remarks hold for $M = \bigcup_J M_j^A = \bigcup_R M_r^B$, the respective partitions of M into A-cosets and B-cosets. The following results are due to these matters.

Proposition 36. *Let* $\mathbf{S}$ *be a skew lattice and* $\{ J > A, B > M \}$ *a skew diamond in* $\mathbf{S}$*. Then,* $[J : A] \leq \omega[J, B]$ *and, dually,* $[M : A] \leq \omega[M, B]$*. Furthermore,*

$$[J : A]|J_i^A \cap J_k^B| = \omega[J, B]$$

and, dually, $[M : A]|M_j^A \cap M_r^B| = \omega[M, B]$*.*

Proof. First observe that the orthogonality of A and B in J is implied by Proposition 17. Thus, due to Proposition 15 and Theorem 3 (i), each B-coset in J intersects all A-cosets in J and no two A-cosets in J intersect, respectively. Hence, $[J : A] \leq \omega[J, B]$.

Fixing $j \in J$, each $y \in B \vee j \vee B$ is in J. Due to Theorem 3 (iii), there exist $a \in A$ such that $y \in A \vee x \vee A$, for some $x \in a \vee J \vee a$. Moreover, B is the union of all these A-cosets in J with x being an element of the transversal $a \vee J \vee a$ and therefore determining a unique coset. Thus,

$$\bigcup_{x \in a \vee J \vee a} (A \vee x \vee A) \cap (B \vee j \vee B) = B \vee j \vee B$$

so that the equality $[J : A]|J_i^A \cap J_k^B| = \omega[J, B]$ holds as required. Similar remarks hold for the dual case regarding A-cosets in M and B-cosets in M. ☐

The result of Proposition 36 permits us to revisit the proofs of the following results, stated by J. Leech, and relating them with the achievements by this innovative approach.

Proposition 37. *[12] Let* $\{ J > A, B > M \}$ *be a skew diamond in a skew lattice* $\mathbf{S}$*. Then,* J *is finite if both* $\omega[J,A]$ *and* $\omega[J,B]$ *are finite, in which case* $|J|$ *divides*

$$\frac{\omega[J,A].\omega[J,B]}{\omega[J,M]}.$$

Dual remarks hold for M*.*

Proof. Assume that $\omega[J,A]$ and $\omega[J,B]$ are finite, By Proposition 35,

$$|J| = [J:A].\omega[J,A].$$

On the other hand, Proposition 36 implies that $[J:A]|J_i^A \cap J_k^B| = \omega[J,B]$. Thus, $[J:A]$ is finite implying the finitude of $|J|$. In that case, again by Proposition 35,

$$\frac{\omega[J,A].\omega[J,B]}{\omega[J,M]} = \frac{|J|^2.[J:M]}{|J|.[J:A].[J:B]} = |J|.\frac{[J:M]}{[J:A].[J:B]}.$$

□

Proposition 38. *[16] Let* $\{ J > A, B > M \}$ *be a skew diamond in a skew lattice* $\mathbf{S}$*. If* J_i^A *and* J_k^B *are finite for any pair of indices* $(j,k) \in I \times K$*, then*

$$\mid J \mid = \frac{\omega[A,J]\omega[B,J]}{|J_i^A \cap J_k^B|}.$$

Similar remarks hold for the cosets of A *and* B *in the meet-class* M*.*

Proof. Assuming the finitude of J_i^A and J_k^B for any pair of indices $(j,k) \in I \times K$, the finitude of J follows from Proposition 37.

The second part of the statement is due to Proposition 36 as $[J:A]|J_i^A \cap J_k^B| = \omega[B,J]$ so that

$$\frac{\omega[A,J].\omega[B,J]}{|J_i^A \cap J_k^B|} = \omega[A,J][J:A] = \mid J \mid.$$

□

In the following result we give a characterization of finite symmetric skew lattices, inspired by communications with J. Leech, that is due to such combinatorial nature.

Proposition 39. *Let* $\{ J > A, B > M \}$ *be a skew diamond in a skew lattice* $\mathbf{S}$*. Then,* $|J_i^A \cap J_k^B| \geq \omega[J,M]$*. Furthermore, if* $\mathbf{S}$ *is symmetric then*

$$|J_i^A \cap J_k^B| = \omega[J,M] \tag{4}$$

and, dually,

$$|M_j^A \cap M_r^B| = \omega[J, M]. \tag{5}$$

Conversely, whenever **S** *is finite, the equalities (4) and (4) imply that the correspondent skew diamond is symmetric.*

Proof. The first two assertions are direct consequences of Proposition 5, respectively. Conversely, assume the finitude of the skew lattice **S** and let $\{ J > A, B > M \}$ be a skew diamond in **S** that satisfies the equalities (4) and (5). Due to Proposition 5 and to the finitude of **S**, for every $m \in M$ and every $j \in J$,

(i) the identity (5) implies that $J \wedge m \wedge J = (A \wedge m \wedge A) \cap (B \wedge m \wedge B)$;

(ii) and the identity (4) implies that $M \vee j \vee M = (A \vee j \vee A) \cap (B \vee j \vee B)$.

Hence, we are in the conditions of Proposition 5 and, therefore, **S** is symmetric. □

Such a characterization is available for normal (and conormal) skew lattices, as a direct consequence of Proposition 26.

Proposition 40. *[18] Let* **S** *be a skew lattice. Then* **S** *is normal if and only if* $[B : A] = 1$ *and* $\omega[A, B] = |B|$ *for each non comparable* $\mathcal{D}$*-classes* $A > B$*. Dually, if* **S** *is conormal if and only if* $[A : B] = 1$ *and* $\omega[A, B] = |A|$ *for each non comparable* $\mathcal{D}$*-classes* $A > B$*.*

We turn to examine the more general case of strictly categorical skew lattices. From it, in conjugation with Propositions 34 and 42 we are able to achieve counting results regarding distributive skew lattices.

Proposition 41. *[18] Given a skew chain* $A > B > C$ *in a skew lattice* **S** *with both* A *and* C *finite, then*

$$|B| \geq \frac{\omega[A, B]\omega[B, C]}{\omega[A, C]}.$$

Furthermore, $A > B > C$ *is strictly categorical if and only if*

$$|B| = \frac{\omega[A, B]\omega[B, C]}{\omega[A, C]} \tag{6}$$

so that in particular B *is also finite. With* B *finite,* $A > B > C$ *is strictly categorical if and only if*

$$|B| = \omega[A, C][B : A][B : C]. \tag{7}$$

Proposition 42. *Let* $A > B > C$ *be a skew chain where* A *and* C *are finite. Then,* $A > B > C$ *is distributive if and only if*

$$|B_i| = \frac{\omega[A, B]\omega[B, C]}{\omega[A, C]}$$

for each AC*-component* B_i *of* B*.*

Corollary 43. *Let $A > B > C$ be a skew chain where A and C are finite. If B has $n < \infty$ AC-components, then $A > B > C$ is distributive if and only if*

$$|B| = n\frac{\omega[A,B]\omega[B,C]}{\omega[A,C]}.$$

Example. *The situation in B is illustrated in the following eggbox figure where rows represent A-cosets in B, columns represent C-cosets in B and their intersections are copies in B of A-cosets in C or C-cosets in A. Here $|B| = 24$ with $\omega[A,B] = 8$, $\omega[B,C] = 6$ and $\omega[A,C] = 2$.*



Acknowledgments

The author would like to thank the challenging suggestions by Jonathan Leech, the insight comments of Karin Cvetko-Vah, and the revealing discussions at Andrej Bauer's seminar.

References

[1] R. Bignall and J. Leech, Skew boolean algebras and discriminator varieties. In *Algebra Universalis* **33** (1995), 387398.

[2] K. Cvetko-Vah. On strongly symmetric skew lattices. *Algebra Universalis* **66** (2011), 99113.

[3] K. Cvetko-Vah, M. Kinyon, J. Leech, and M. Spinks. Cancellation in skew lattices. *Order* **28** (2011), 932.

[4] K. Cvetko-Vah and J. Pita Costa. On the coset laws for skew lattices in rings. *Novi Sad J. Math* **40** (2010), 1125.

[5] K. Cvetko-Vah and J. Pita Costa. On the coset laws for skew lattices. *Semigroup Forum* **83** (2011), 395411.

[6] K. Cvetko-Vak. Skew lattices in rings. *PhD thesis.* University of Ljubljana, 2005.

[7] J. A. Green. On the structure of semigroups. *Annals of Mathematics* **54** (1951), 163172.

[8] J.M. Howie. *An introduction to semigroup theory.* Academic Press (1976).

[9] M. Kinyon and J. Leech. Categorical skew lattices. *submitted* (2012).

[10] M. Kinyon and J. Leech. Distributivity in skew lattices. *preprint* (2012).

[11] J. Leech. Towards a theory of noncommutative lattices. *Semigroup Forum* **34** (1986),117120.

[12] J. Leech. Skew lattices in rings. *Algebra Universalis* **26** (1989), 4872.

[13] J. Leech. Skew boolean algebras. *Algebra Universalis* **27** (1990), 497506.

[14] J. Leech. Normal skew lattices. *Semigroup Forum* **44** (1992), 1–8.

[15] J. Leech. The geometric structure of skew lattices. *Trans. Amer. Math. Soc.* **335** (1993), 823842.

[16] J. Leech. Recent developments in the theory of skew lattices. *Semigroup Forum* **52** (1996), 7-24.

[17] J. Pita Costa, On the coset structure of a skew lattice. *Demonstratio Math.* **44** (2011), 673–692.

[18] J. Pita Costa, Coset laws for categorical skew lattices. *Algebra Univers.* DOI 10.1007/s00012-012-0194-z (2012). **68** (2012), 75-89.

[19] J. Pita Costa, *On the coset structure of skew lattices.* Doctoral Thesis, University of Ljubljana (2012).

[20] J. Pita Costa, On the coset structure of a skew lattice. *Demonstratio Math.* submmitted (2012).

[21] B. M. Schein. Pseudosemilattices and pseudolattices. *Amer. Math. Soc. Transl., 119* (1983), 116.

[22] M. Spinks. On middle distributivity for skew lattices. *Semigroup Forum* **61** (2000), 341345.